

AgriGuard Fluroxypyr

MAPP 13335

This product is an emulsifiable concentrate containing 200g/l (20.4% w/w) fluroxypyr (acid equivalent) present as 288g/L of the 1-methyl heptyl ester, and aromatic naphtha.

AgriGuard Fluroxypyr is a post-emergence herbicide for the control of a range of annual and perennial weeds in winter and Spring Cereals, forage maize and Grassland.



VERY TOXIC TO AQUATIC ORGANISMS
**MAY CAUSE LONG TERM
ADVERSE EFFECTS** IN THE
AQUATIC ENVIRONMENT

**DANGEROUS FOR
THE ENVIRONMENT**



FLAMMABLE
IRRITATING TO EYES, SKIN AND
RESPIRATORY SYSTEM

HARMFUL MAY CAUSE LUNG DAMAGE IF
SWALLOWED

HARMFUL

**MAY CAUSE
SENSITISATION** BY SKIN CONTACT
**VAPOURS MAY CAUSE
DROWSINESS OR
DIZZINESS**

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/situations	Maximum Individual dose (litres product per hectare)	Maximum Total dose (litres product per hectare, per crop)	Latest time of application
Wheat (winter), Barley (winter)	2	2	Before flag leaf sheath opening stage
Oats (winter), rye, Triticale, durum wheat	1	1	Before second node detectable stage
Wheat (spring), Barley (spring)	0.75	0.75	Before flag leaf sheath extending stage
Oats (spring)	0.75	0.75	Before second node detectable stage
Forage Maize	1	1	Before seven leaves unfolded stage
Grassland	2	2 litres product per hectare, per year	-

Other Specific Restrictions:

1. When the product is applied as a spot treatment to permanent grassland, rotational grass and newly sown grass leys, the maximum concentration must not exceed 30ml product per 10 litres water.
2. A maximum total dose of 0.75 litres per hectare must be observed for applications made to cereals between crop emergence in the year of planting and 1st February in the year of harvest.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

Engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

IF SWALLOWED do not induce vomiting: seek medical advice immediately and show this container or label.

AVOID CONTACT WITH SKIN.

IN CASE OF CONTACT WITH EYES, rinse immediately with plenty of water and seek medical advice.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show label where possible).

Environmental Protection

DO NOT CONTAMINATE WATER with product or its container. Do not clean application equipment near surface water. Avoid contamination from farmyards and roads.

KEEP ALL LIVESTOCK out of treated areas for at least 3 days and until the foliage of poisonous weeds such as ragwort have died and become unpalatable.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank, and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN

To avoid risks to man and the environment, comply with the instructions for use.

This material and its container must be disposed of in a safe way.

Use appropriate containment to avoid environmental contamination

Transport Information:

UN 1993

ADR/RID Class: 3

Packaging Group: III

Net Contents: 5 Litres 

PROTECT FROM FROST

SHAKE WELL BEFORE USE

Marketing Company: AgriGuard Limited, Units 1& 2, Block B, Broomfield Business Park, Malahide, Co. Dublin, Ireland.

Tel: +353-1-8462044 Fax: +353-1-8462489

www.agriguard.ie

Code: R356/UK/02

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

AGRIGUARD FLUROXYPYR is a selective post-emergence herbicide for the control of a range of annual and perennial weeds in Cereals, Forage Maize and Grassland.

RESTRICTIONS

Do not treat crops that are or have been under stress from any cause, including poor soil or cultural conditions, adverse climatic conditions, waterlogging or drought, pest or disease attack, nutrient deficiency, or any other cause.

For optimum results, apply **AGRIGUARD FLUROXYPYR** when the weeds are small and actively growing. The crop should be growing strongly to provide good competition with the weeds. This is particularly important with cleavers especially where HBN's are not included in the tank mix. HBN's are products containing ioxynil and/or bromoxynil (refer to Compatibility section).

Crops which are undersown with grass may be sprayed provided the grasses are tillering.

Do not apply **AGRIGUARD FLUROXYPYR** to crops undersown with clover or other legume-containing mixtures.

Keep livestock out of treated areas for at least 3 days and until poisonous weeds such as ragwort have died and become unpalatable.

Avoid spray drift onto other crops or areas.

Do not spray if frost is likely to occur soon after spraying, during periods of frosty weather, or during periods of low night temperatures.

Do not roll or harrow 1 week before or after application.

Apply to dry foliage, when rain is not imminent.

Contract agents should be consulted before using on crops grown for seed.

AGRIGUARD FLUROXYPYR will not control weeds that emerge after application.

ONLY one application of **AGRIGUARD FLUROXYPYR** may be made per crop or year (see statutory conditions of approval).

When the product is applied as a spot treatment to permanent grassland, rotational grass and newly sown grass leys the maximum concentration must not exceed 30ml product per 10 litres water.

A maximum total dose of 0.75 litres per hectare must be observed for applications made to cereals between crop emergence in the year of planting and 1st February in the year of harvest.

CROP SPECIFIC INFORMATION

Grassland

Do not treat crops where clover or other legumes are an important part of the sward.

Overall Treatment:

AGRIGUARD FLUROXYPYR is applied at two rates; 2.0 litres/ha on established grassland (in 300 – 400 litres of water/ha), and 0.75 litres/ha on newly sown grass leys (in 200-400 litres water/ha). Always use the higher volume of water recommended when the grass growth is thick, or weeds are at the maximum recommended growth stage for treatment. Always use a water volume that will ensure good spray coverage of the weeds.

Spot treatment:

Where weed populations are too low to justify overall spraying, **AGRIGUARD FLUROXYPYR** may be applied as a spot treatment to established grassland and newly sown grass leys, using a spray concentration of 30ml **AGRIGUARD FLUROXYPYR** in 10 litres of water. Spray to evenly wet the foliage, to just before the point of run-off.

Keep livestock out of treated areas until the foliage of any poisonous weeds such as Ragwort has died and become unpalatable; under certain conditions, this may take several weeks. The grazing / cutting management regime must be planned in advance to allow sufficient time for this to occur.

A. Established Grassland

AGRIGUARD FLUROXYPYR is recommended for the control of certain perennial weeds in established grassland.

Apply 2.0 litres of **AGRIGUARD FLUROXYPYR** per hectare in 300 – 400 litres of water per hectare, according to the optimum timing shown below. Do not apply more than a total maximum dose of 2.0 litres per hectare per year.

Weed Species	Timing and Notes
Curled Dock	After cutting or grazing, ensure that sufficient weed foliage is present and that the Docks are actively growing before spraying. Delay spraying for 2 – 3 weeks after cutting or grazing. Apply in Spring when the Docks are at the rosette stage, and 15 – 20 cm high. A follow-up treatment in Spring of the following year may be required where dock populations are dense and/or well established.
Broad-leaved Dock	
Dandelion	Apply in the Spring when growing actively, but prior to flowering.
Common Nettle	For a reduction in top growth, apply in the Spring when growing actively, but prior to flowering (normally up to mid June).
Bramble, Broom	Spray when the plant is actively growing in June-August but before growth begins to slow in the autumn. Incomplete eradication may result if the foliage is inadequately wetted.

B. Newly-sown grass leys (Autumn and Spring application)

AGRIGUARD FLUROXYPYR is recommended for the control of Common Chickweed in newly-sown grass leys. The grass must have at least 3 fully expanded leaves at the time of treatment. Common Chickweed will be controlled at up to 5 cm.

Apply 0.75 litres per hectare of **AGRIGUARD FLUROXYPYR** in 200 – 400 litres of water per hectare. Apply in the autumn or spring when the weeds are actively growing. Do not apply more than a total maximum dose of 0.75 litres of **AGRIGUARD FLUROXYPYR** per hectare per year.

Forage Maize

AGRIGUARD FLUROXYPYR is recommended for the control of Black Nightshade from cotyledons up to 6 leaves stage.

AGRIGUARD FLUROXYPYR will not control Black Nightshade which has not emerged at the time of application.

The maize must be between the 3 – 6 leaf stage and must not be over 20 cm in height.

Do not apply AGRIGUARD FLUROXYPYR if the side roots have started to develop on the first node.

Apply 1.0 litre per hectare of **AGRIGUARD FLUROXYPYR** in 200 – 300 litres of water per hectare. Do not apply more than a total maximum dose of 1.0 litre of **AGRIGUARD FLUROXYPYR** per hectare per year.

Do not apply **AGRIGUARD FLUROXYPYR** in a tank mix with any other product or if the crop is beyond the recommended growth stage. Boom overlap must be avoided.

Cereals

A. Spring Application to Winter and Spring Cereals

AGRIGUARD FLUROXYPYR may be applied in the Spring according to the Crops, Timing and Rates listed in the table below.

The recommended spray volume is 150-400 litres of water per hectare. Only use 150 litres of water per hectare on small weeds in very open crops. Increase the volume to 400 litres of water per hectare in thick crops and where weed cover is thick.

Crop	Timing of Application	Rate L/ha	Weeds Controlled & Maximum Growth Stage for Control
Winter Wheat Winter Barley	Apply from the 2 leaf stage (GS 12) until before flag leaf sheath opening stage (GS 45)	1.0	Cleavers (flowering) Common Chickweed (flowering) Common hemp-nettle (flowering) Field forget-me-not (flowering) Black Bindweed (6 leaves) Henbit dead nettle (4 leaves) Red dead-nettle (4 leaves) Knotgrass (2 leaves)
Winter Oats Winter Rye Durum Wheat Triticale	Apply from the 2 leaf stage (GS 12) until before the second node detectable stage (GS 31)	1.0	Common Fumitory (2 leaves) Groundsel (checked at 2 leaves) Mayweeds (checked at 2 leaves) Redshank (checked at 2 leaves) Common field-speedwell (checked at 2 leaves) Ivy-leaved Speedwell (checked at 2 leaves) Pale persicaria (checked at 2 leaves)
Spring Wheat Spring Barley	Apply from the 2 leaf stage (GS 12) up to and including the flag leaf ligule just visible stage (GS39).	0.75	Cleavers (10 cm) Common Chickweed (10 cm) Common hemp-nettle (10 cm) Field forget-me-not (5 cm) Black Bindweed (4 leaves) Knotgrass (2 leaves) Common Fumitory (2 leaves) Corn spurrey (2 leaves)
Spring Oats	Apply from the 2 leaf stage (GS 12) until before second node detectable stage (GS31).	0.75	Groundsel (checked at 2 leaves) Mayweeds (checked at 2 leaves) Redshank (checked at 2 leaves) Common field-speedwell (checked at 2 leaves) Ivy-leaved Speedwell (checked at 2 leaves) Pale persicaria (checked at 2 leaves)

Cereal crops undersown with grass may be treated if the grass has reached the tillering stage. Do not treat crops undersown with clover or other legumes.

B. Spring Application in Tank Mix with HBNs*

AGRIGUARD FLUROXYPYR can be tank mixed with HBNs for use in crops of winter and spring cereals in the Spring according to the Crops, Timing and Rates listed in the tables below.

Do not spray if frost is imminent. Application to winter oats should be delayed until risk of frost is over.

LOW RATE MIXTURE

(recommended where cleavers, chickweed, hemp-nettle and forget-me-not are relatively small; see maximum growth stages below)

Crop	Timing of Application	Rate L/ha	Weeds Controlled & Maximum Growth Stage for Control
Winter Wheat Winter Barley	Apply from the 2 leaf stage (GS 12) until before the second node detectable stage (GS 31)	0.75 + recommended rate of HBN	Cleavers (25 cm) Common Chickweed (flowering) Common Hemp-nettle (15 cm) Field Forget-me-not (10 cm) Black Bindweed (10 cm) Henbit Dead-nettle (10 cm) Red Dead-nettle (10 cm) Knotgrass (10 cm) Common Fumitory (10 cm)
Winter Oats Winter Rye Durum Wheat	Apply from the 2 leaf stage (GS 12) until before the second node detectable stage (GS 31)	0.75 + recommended rate of HBN	Groundsel (Up to 5 cm) } Mayweeds (Up to 5 cm) } Redshank (Up to 5 cm) } depends on type and Common Field-speedwell (Up to 5 cm) } rate of HBN used* Ivy-leaved Speedwell (Up to 5 cm) } Pale Persicaria (Up to 5 cm) } Bugloss (Up to 6 true leaves) } Charlock (Up to 6 true leaves) } Common Poppy (Up to 6 true leaves) } Corn Marigold (Up to 6 true leaves) } Fat-hen (Up to 6 true leaves) } Field Penny-cress (Up to 6 true leaves) } Scarlet Pimpernel (Up to 6 true leaves) } Shepherd's Purse (Up to 6 true leaves) } Small Nettle (Up to 6 true leaves) } Volunteer Rape (Up to 6 true leaves) } Wild Radish (Up to 6 true leaves) }

Spring Wheat Spring Barley Spring Oats	Apply from the 2 leaf stage (GS 12) until before second node detectable stage (GS31).	0.5 + recommended rate of HBN	Cleavers (5 cm) Common Chickweed (10 cm) Common Hemp-nettle (10 cm) Field Forget-me-not (5 cm) Black Bindweed (5 cm) Knotgrass (5 cm) Common Fumitory (5 cm) Corn Spurrey (Up to 6 true leaves) } Groundsel (Up to 6 true leaves) } Mayweeds (Up to 6 true leaves) } depends on type and Redshank (Up to 6 true leaves) } rate of HBN used* Common Field-speedwell (Up to 6 true leaves) } Ivy-leaved Speedwell (Up to 6 true leaves) } Pale Persicaria (Up to 6 true leaves) } Fat-hen (Up to 6 true leaves) } Common Orache (Up to 6 true leaves) } Common Poppy (Up to 6 true leaves) } Shepherd's Purse (Up to 6 true leaves) }
--	---	-------------------------------	--

* HBNs are products containing ioxynil and/or bromoxynil (refer to Compatibility section). HBN tank mix partners recommended are MAFF/MAPP 09470[†], 10005[°].

[†] For use only in barley, oats, rye, triticale and wheat.

[°] For use only in barley, oats, rye and triticale.

Use tank mix partners in accordance with the manufacturers instructions.

FULL RATE MIXTURE

(recommended where cleavers, chickweed, hemp-nettle and forget-me-not are more advanced; see maximum growth stages below)

Crop	Timing of Application	Rate L/ha	Weeds Controlled & Maximum Growth Stage for Control
Winter Wheat Winter Barley	Apply from the 2 leaf stage (GS 12) until before the second node detectable stage (GS 31)	1.0 + recommended rate of HBN	Cleavers (flowering) Common Chickweed (flowering) Common Hemp-nettle (flowering) Field Forget-me-not (flowering) Black Bindweed (10 cm) Henbit Dead-nettle (10 cm) Red Dead-nettle (10 cm) Knotgrass (10 cm) Common Fumitory (10 cm)

Winter Oats Winter Rye Durum Wheat Triticale	Apply from the 2 leaf stage (GS 12) until before the second node detectable stage (GS 31)	1.0 + recommended rate of HBN	Groundsel (Up to 5 cm) } Mayweeds (Up to 5 cm) } Redshank (Up to 5 cm) } depends on type and Common Field-speedwell (Up to 5 cm) } rate of HBN used* Ivy-leaved Speedwell (Up to 5 cm) } Pale Persicaria (Up to 5 cm) } Bugloss (Up to 6 true leaves) } Charlock (Up to 6 true leaves) } Common Poppy (Up to 6 true leaves) } Corn Marigold (Up to 6 true leaves) } Fat-hen (Up to 6 true leaves) } Field penny-cress (Up to 6 true leaves) } Scarlet Pimpernel (Up to 6 true leaves) } Shepherd's Purse (Up to 6 true leaves) } Small Nettle (Up to 6 true leaves) } Volunteer Rape (Up to 6 true leaves) } Wild Radish (Up to 6 true leaves) }
Spring Wheat Spring Barley Spring Oats	Apply from the 2 leaf stage (GS 12) until before second node detectable stage (GS31).	0.75 + recommended rate of HBN	Cleavers (15 cm) Common Chickweed (flowering) Common Hemp-nettle (15 cm) Field Forget-me-not (10 cm) Black Bindweed (10 cm) Knotgrass (10 cm) Common Fumitory (10 cm) Corn Spurrey (Up to 6 true leaves) } Groundsel (Up to 6 true leaves) } Mayweeds (Up to 6 true leaves) } depends on type and Redshank (Up to 6 true leaves) } rate of HBN used* Common Field-speedwell (Up to 6 true leaves) } Ivy-leaved Speedwell (Up to 6 true leaves) } Pale Persicaria (Up to 6 true leaves) } Fat-hen (Up to 6 true leaves) } Common Orache (Up to 6 true leaves) } Common Poppy (Up to 6 true leaves) } Shepherd's Purse (Up to 6 true leaves) }

* HBNS are products containing ioxynil and/or bromoxynil (refer to Compatibility section). HBN tank mix partners recommended are MAFF/MAPP 09470[†], 10005[°].

[†] For use only in barley, oats, rye, triticale and wheat.

[°] For use only in barley, oats, rye and triticale.

Use tank mix partners in accordance with the manufacturers instructions.

Note: In spring cereals the full rate mixture will also give control of black bindweed, knotgrass and fumitory at later growth stages than with the low rate mixture (see tables for details of maximum growth stages controlled).

C. Autumn Application to Winter Wheat and Winter Barley

AGRIGUARD FLUROXYPYR may be applied in the Autumn to winter wheat and winter barley only according to the Crops, Timing and Rates listed in the table below.

Weeds which germinate after spraying will not be controlled by an Autumn application of **AGRIGUARD FLUROXYPYR**.

For optimum results, weeds must be actively growing and the crop competitive. This is particularly important with cleavers.

Do not spray if frost is imminent.

Crop	Timing of Application	Rate* L/ha	Weeds Controlled & Maximum Growth Stage for Control
Winter Wheat Winter Barley	Apply from the 2 leaf stage (GS 12) up until the end of February, or first node detectable stage (GS 31), whichever is sooner. Spring Application recommendations apply from March onwards	0.5-0.75 + recommended rate of HBN	Cleavers (5 cm) Common Chickweed (5 cm) Field Forget-me-not (5 cm) Henbit Dead-nettle (6 true leaves) Red Dead-nettle (6 true leaves) Charlock (6 true leaves) } Common Poppy (6 true leaves) } Groundsel (6 true leaves) } Mayweeds (6 true leaves) } Shepherd's Purse (6 true leaves) } depends on type and Common Field-speedwell (6 true leaves) } rate of HBN used Ivy-leaved Speedwell (6 true leaves) } Volunteer rape (6 true leaves) }

* The 0.75l/ha rate of **AGRIGUARD FLUROXYPYR** generally gives more consistent results particularly if growing conditions are not ideal. This rate is essential for the control of cleavers.

HBN's are products containing ioxynil and/or bromoxynil (refer to Compatibility section). HBN tank mix partners recommended are MAFF/MAPP 09470, and 10005. Use in accordance with the manufacturers instructions.

D. Volunteer potato control in winter wheat and winter barley.

Potato shoots from 10 cm to 40 cm in height can be controlled by application of 2.0 litres **AGRIGUARD FLUROXYPYR** in 300 to 400 litres of water per hectare when the crop is between the third node detectable and flag leaf ligule just visible growth stage (GS 33 to GS 39 inclusive).

NOTE

AGRIGUARD FLUROXYPYR will not completely control top growth but will check the potato plants sufficiently to reduce their competitiveness. There will also be a useful reduction in viability of daughter tubers.

AGRIGUARD FLUROXYPYR will only control shoots emerged at the time of spraying and it is therefore best used later in its timing window.

AGRIGUARD FLUROXYPYR will only be effective if there is adequate foliage on actively growing potato shoots to absorb the herbicide and good coverage is obtained, hence the need for higher water volumes.

IMPORTANT

When **AGRIGUARD FLUROXYPYR** is used at the 2.0 litre / ha rate, the following extra warnings apply:

- Avoid overlapping spray bouts
- **Do not** tank mix **AGRIGUARD FLUROXYPYR** with any other product when used at the 2.0 l/ha rate.
- Do not plant field beans, peas or other legumes in the same year as treatment, nor in the spring of the year following treatment.
- Any straw from crops treated with **AGRIGUARD FLUROXYPYR** at 2.0 litres per hectare must only be used for animal bedding and not incorporated back into the soil.

Manure from the animal bedding must only be returned to grassland or land to be cropped with cereals.

GENERAL INFORMATION

Mixing Instructions

Shake the container before use. Place half the required amount of clean water in the spray tank and commence agitation. Add the required amount of **AGRIGUARD FLUROXYPYR**. Rinse all used containers thoroughly and add the washings to the spray tank. The use of sprayer mounted pressure rinsing equipment is advised. If not available, containers should be manually rinsed three times. Add the remaining water requirement and continue agitation. The spray mix must be used immediately, maintaining continuous agitation until spraying is completed. Do not allow the spray mixture to stand. Immediately after use wash sprayer and other equipment thoroughly with water and detergent. Traces of **AGRIGUARD FLUROXYPYR** in the spray tank could damage susceptible crops when the sprayer is used subsequently. Dispose of empty rinsed containers according to the MAPP/HSE Code of Practice for the Safe Use of Pesticides on Farms and Holdings.

Method of Application

Apply by means of a conventional hydraulic nozzle tractor-mounted or trailed sprayer (with or without air-assistance). Ensure that the sprayer is correctly calibrated to apply the recommended water volume and spray quality and the boom height and nozzles are adjusted to give even coverage of the target weeds.

In grassland, small areas may be treated by knapsack or other hand-operated sprayer. Spray individual weeds to the point just prior to run-off. Ensure that the sprayer is correctly calibrated to apply the recommended dose and water volume.

Water Volume

Use a volume of water within the recommended range that will give a good even cover of the weed foliage. See recommendations under Crop Specific Information. The minimum recommended pressure is 2 to 3 bars.

The lowest volume of water should only be used in very open crops on small weeds. On later applications where the crop is dense, the spray volume should be increased to 400 litres of water per hectare.

Spray Quality

Apply as a MEDIUM spray (as defined by BCPC).

Following Crops and Residues from Treated Crops

Do not plant beans or other legumes in the same year as an application of **AGRIGUARD FLUROXYPYR** at the highest rate or total maximum dose applied of 2.0 litres per hectare. Do not plant peas or other legumes in the spring following an application of **AGRIGUARD FLUROXYPYR** at the highest rate or total maximum dose applied of 2.0 litres per hectare

Any straw from crops treated with **AGRIGUARD FLUROXYPYR** at 2.0 litres per hectare must only be used for animal bedding and not incorporated back into the soil. Manure from the animal bedding must only be returned to grassland or land to be cropped with cereals.

Tank Mixes

To broaden the range of weeds controlled, **AGRIGUARD FLUROXYPYR** can be tank mixed with a number of herbicides. If using tank mixes, unless directed otherwise the preferred order of addition of products to the tank is: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product must be added to a half-full sprayer and be fully dispersed before the addition of the next product.

Tank mixes must only be applied within the label recommendations of every product in the mix.

Compatibility

AGRIGUARD FLUROXYPYR is compatible with a wide range of products. The weed spectrum can be broadened by tank-mixing with a range of different herbicides.

AGRIGUARD FLUROXYPYR is compatible with a number of HBN products. HBNs are products containing ioxynil and/or bromoxynil. HBN tank mix partners recommended are MAFF/MAPP 09470, and 10005. Use in accordance with the manufacturers instructions.

Contact AgriGuard Ltd for compatibility information on specific tank mixes. Manufacturer's instructions must be followed for each tank-mix component.

Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.